



The Role of Tlrs Signaling in Fungal Infections

Mehdi Taghavi

University of Tehran, Iran.

Abstract:

Fungi are the most common microorganisms found in the environment. Fungal immunity is vital in the treatment of immunocompromised patients, especially those involved with HIV infection, cancer, or organ transplant recipients. There are numerous immune mechanisms against fungal infections, ranging from nonspecific protective mechanisms (innate immunity) to sophisticated acquired one that are activated during an infection or a disease (adaptive immunity) with the former (innate immunity) being the first line of defense against many fungal infections. Pattern recognition receptors (PRRs) such as Toll-like receptors, C-type lectin receptors (CLRs), and galectin family proteins exist on the surface of host cells to identify PAMPs in fungi. The major components of fungal cell wall are beta-glucan and chitin. Depending on the molecular size, chitin induces innate or adoptive immunity activity while beta-glucan activates innate immunity during fungal infections. PRRs, existing on different cells such as phagocytes, epithelial, and dendritic cells initiate intracellular events that activate immune system and fungal clearance. TLRs are pathogen recognition receptors that detect specific PAMPs in microbes. They exist on host cell surfaces or into cytoplasm regions, identifying microbial ligands. The interaction between TLRs and PAMPs activates immune cells, followed by antimicrobial proteins and cytokines secretion, respiratory burst, and phagocyte degranulation. TLRs can be divided into two groups: those like TLR1, 2, 4, and 5 that exist on the surface of plasma membrane and TLR3, 7, 8, and 9 which are placed in intracellular components such as endosomes. The type of immunity provoked depends on the cells involved, e.g., neutrophils, macrophages, and some other cell types generally promote innate immune



responses, while antigen uptake by dendritic cells results in differentiation of naïve T cells and specific immunity responses. This paper describes the role of Tolllike receptors for the identification of immune responses to fungal infections.

Biography:

Mehdi Taghavi ia a PHD Student at University of Tehran from Iran.

Publication of speakers:

1. Honarvar, Behnam & Lankarani, Kamran & Taghavi, Mehdi & Vahedi, Ghasem & Mortaz, Esmail. (2019). Biomarker-guided antifungal stewardship policies for patients with invasive candidiasis. *Current Medical Mycology*. 4. 10.18502/cmm.4.4.385.
2. Taghavi, Mehdi & Mortaz, Esmail & khosravi, Alireza & Folkerts, Gert & Vahedi, Ghasem & Varahram, Mohammad & Kazempour- Dizaji, Mehdi & Garssen, J. & Adcock, Ian. (2018). Zymosan attenuates melanoma growth progression, increases splenocyte proliferation and induces TLR2 and TNF- α expression in mice. *Journal of inflammation*. 15. 10.1186/s12950-018-0182-y.
3. Taghavi, Mehdi & Khosravi, Ali Reza & Mortaz, Esmail & Nikaein, Donya & Athari, Seyyed. (2016). Role of pathogen-associated molecular patterns (PAMPS) in immune responses to fungal infections. *European Journal of Pharmacology*. 808. 10.1016/j.ejphar.2016.11.013.

[International Conference on New Frontiers in Clinical Microbiology and Infectious Diseases | November 19-20, 2020 | London, UK](#)

Citation: Mehdi Taghavi; The Role of Tlrs Signaling in Fungal Infections; Euro Microbiology 2020; November 19-20, 2020; London, UK